

# Eclipsing binaries in ASAS catalog

B. Paczyński<sup>1\*</sup> and D. Szczygiel<sup>2</sup>, B. Pilecki<sup>2</sup>, G. Pojmański<sup>2</sup> †

<sup>1</sup>*Princeton University Observatory, Peyton Hall, Princeton, NJ 08544, USA*

<sup>2</sup>*Warsaw University Observatory, Al. Ujazdowskie 4, PL-00-478, Poland*

Accepted –. Received – ; in original form –

## ABSTRACT

ASAS is a long term project to monitor bright variable stars over the whole sky. It has discovered 50,122 variables brighter than  $V < 14$  mag south of declination  $+28^\circ$ , and among them 11,099 eclipsing binaries. We present a preliminary analysis of 5,384 contact, 2,957 semi-detached, and 2,758 detached systems. The statistics of the distribution provides a qualitative confirmation of decades old idea of Flannery and Lucy that W UMa type binaries evolve through a series of relaxation oscillations: ASAS finds comparable number of contact and semidetached systems.

The most surprising result is a very small number of detached eclipsing binaries with periods  $P < 1$  day, the systems believed to be the progenitors of W UMa stars. As many (perhaps all) contact binaries have companions, there is a possibility that some were formed in a Kozai cycle, as suggested by Eggleton and his associates.

**Key words:** stars: eclipsing – stars: binary – stars: evolution

## 1 INTRODUCTION TO ASAS

ASAS - All Sky Automated Survey, is a long term project dedicated to detection and monitoring variability of bright stars. This paper presents the results of several years of observations done at the Las Campanas Observatory with a single instrument: a telescope with the aperture of 7 cm, the focal length of 20 cm, done through a standard V-band filter and a  $2K \times 2K$  CCD camera with  $15\mu$  pixels from Apogee (Pojmański 1997, 1998, 2000, 2002, 2003, Pojmański and Maciejewski 2004, 2005, Pojmański, Pilecki and Szczygiel 2005). More information about ASAS is provided on the WWW:

<http://www.astrouw.edu.pl/~gp/asas/asas.html>

<http://archive.princeton.edu/~asas/>

The variable stars were discovered quasi - uniformly for declination  $< +28^\circ$ , covering almost  $3/4$  of the full sky. Fig. 1 shows the distribution of ASAS variables in the sky in the Galactic coordinates. The Milky Way is clearly visible, together with the dust lanes. The distribution of ASAS stars as a function of V-mag is shown on Fig. 2 for all stars (17,000,000), all variable stars (50,122), and all eclipsing binaries, the latter divided into Eclipsing Contact binaries (EC, 5,384), Eclipsing Semi Detached binaries (ESD, 2,957), and Eclipsing Detached binaries (ED, 2758). The statistics for stars with  $8 < V < 12$  mag appears to be approximately complete, but the efficiency falls rapidly within the range of  $12 < V < 14$  mag as the detection limit is approached. Also,

the statistics deteriorates for  $8 < V$  because of saturation effects.


All stars were observed for about 5 years, with a small subset for 8 years. Typical number of photometric measurements was several hundred. The distribution of this number is shown in Fig. 3. The total number of photometric measurements was 2,916,000. As ASAS continues its operation the number of measurements will increase, approximately 100 V-band photometric measurements per year per variable. We intend to continue the project indefinitely, with some upgrades. While only V-band results are reported here, the I-band photometry was accumulated, and several years of data are already stored on a RAID-5 disk system. However, it will take another year to process I-band data, well over a Tera-byte. There are more stars detectable in the I-band, so we expect that the number of variables will more than double. We are also planning an expansion of ASAS to the northern hemisphere, to fully cover the whole sky.

The results presented in this paper are not final in any sense. Please note, that all data are public domain. A different classification of binaries can be readily done by whoever feels like verifying and/or correcting our presentation.


This is an observational paper, but section 2 gives a short introduction to the main ideas about the structure and evolution of contact binaries. Section 3 gives the information about ASAS data, the classification scheme, and some examples of a diversity of light curves. Section 4 provides simple statistics of ASAS binaries. Finally, in section 5 we make a somewhat speculative discussion based on this observational paper.

\* e-mail: bp@astro.princeton.edu

† e-mail: dszczyg, pilecki, gp@astrouw.edu.pl



**Figure 1.** The distribution of 50,122 ASAS variables in the sky in Galactic coordinates. The Milky Way is clearly seen, as well as the patches of interstellar extinction. The equator is shown with a dashed line. The distribution is limited to declination  $< +28^\circ$ .




**Figure 2.** The distribution of ASAS stars as a function of V-band magnitude. The total number of stars is about 17,000,000. The total number of variable stars is 50,122. The total number of eclipsing binaries is 5,384 for contact systems (EC), 2,957 for semi-detached systems (ESD), and 2,758 for detached systems (ED). Notice, that the efficiency of discovering variable stars declines for  $V > 12$  because the detection limit is approached, and for  $V < 8$  because of saturation effects.

## 2 INTRODUCTION TO CONTACT BINARIES

Contact binaries, also known as W UMa stars, are in a physical contact, with continuously changing brightness because of large tidal distortion of the two components.

The first theoretical milestone in the understanding of contact binaries was due to Lucy (1968a,b), who proposed



**Figure 3.** The histogram of the number of photometric measurements obtained during 5 years of ASAS life. A small subset of data extends back to 8 years.

that the two components share common envelope with the same entropy, thereby making the effective temperature almost constant over the surface of the two stars. As contact binaries have the mass ratio distinctly different from one, most nuclear energy is generated in the more massive component and it is redistributed around the whole surface through a moderately thick convective envelope.

The second theoretical milestone was the recognition of the consequences of the fact that the mass - radius relation for Zero Age Main Sequence (ZAMS) stars is much steeper than for the two Roche lobes. There can be no

stable equilibrium between the two stars with a common envelope. The system evolves through a sequence of relaxation oscillations, with the mass flowing from star A to B, next from B to A, etc. (Flannery 1976, Lucy 1976, Robertson and Eggleton 1977). The cycle repeats on a thermal (Kelvin-Helmholtz) time scale. According to thermal relaxation model the binary oscillates between thermal contact, with the two eclipses of almost equal depth, and a semidetached phase in which one eclipse is much deeper than another.

Hazlehurst (1970) suggested that nuclear evolution of the primary component of a contact binary affects its structure. Stepień (2003, 2005) suggested that the currently more massive primary was originally the less massive of the two. The nuclear burning formed a small helium core, the star expanded and transferred mass to the original secondary. In analogy with Algol systems the currently more massive component is the less evolved, while the present secondary has a small helium core, and it is more advanced in its nuclear evolution.

It is interesting that there is a controversy about thermal relaxation oscillations in W UMa systems. Some authors claim there are no such oscillations (Webbink 2003), while others claim that such oscillations exist (Quin 2003, Yakut and Eggleton 2005, and references therein). ASAS statistics resolves this controversy on purely observational grounds.

### 3 ASAS DATA


Close binaries with a deep common envelope are in thermal contact and they have eclipses of almost equal depth. If the contact is shallow, or if there is no physical contact, then the effective temperatures of the two stars are different, and the two eclipses have different depth.

Theoretical models of relaxation oscillations indicate that the radii of the two components change relatively little throughout the cycle (Flannery 1976, Lucy 1976, Robertson and Eggleton 1977, Yakut and Eggleton 2005, and references therein). With the geometry of the two stars almost unchanged, tidal distortions due to geometry remain almost the same, and the most profound difference in the light curve is the relative depth of the two eclipses.

For the purpose of this paper the classification of eclipsing binaries was done by decomposing their light curves into Fourier coefficients:  $a_1, a_2, a_3, a_4$ , following Ruciński (1997a,b, 1998, 2002) and Pojmański (2002). The classification is shown in Fig. 4 in the  $a_2 - a_4$  plane, with the EC, ESD, and ED binaries shown with different symbols: ESD stars have smaller marks than EC and ED. Some examples of bright binaries of EC, ESD, and ED type are shown in Fig. 5. The parameters of these 15 binaries are listed in Table 1. The type of an eclipsing binary, EC, ESD, or ED, is indicated at the upper right corner of the light curve. The complete version of Table 1, for all eclipsing binaries used in this paper, and with all eight Fourier parameters, is in the file `Fourier.E` on the web page:

[http://www.astrouw.edu.pl/~gp/asas/asas\\_paper\\_data.html](http://www.astrouw.edu.pl/~gp/asas/asas_paper_data.html)

[http://archive.princeton.edu/~asas/asas\\_paper\\_data.html](http://archive.princeton.edu/~asas/asas_paper_data.html) where all the data on which this paper is based are available electronically. These include all photometric measurements




**Figure 4.** Classification of eclipsing binaries in the Fourier coefficients plane  $a_2 - a_4$ . The three symbols refer to contact (EC), semidetached (ESD), and detached (ED) binaries, following Pojmański (2002). Note: when the amplitude of variability is very small the classification is very uncertain, as in the upper right hand corner.

**Table 1.** The most important Fourier parameters for light curves from Fig. 5 are listed along with orbital periods. Each row is separated with a horizontal line. The  $a_2$  parameter can be easily translated to the amplitude,  $a_1$  and  $a_3$  determine the difference between the two eclipses, and  $b_1$  is related to the difference between the two maxima. Full Table 1 is given electronically as file `Fourier.E` on our web page.

ASAS ID	P[days]	$a_1$	$a_2$	$a_3$	$a_4$	$b_1$
125340-5010.6	0.4047	-0.003	-0.158	-0.003	-0.028	-0.003
111621-5427.2	0.4971	-0.026	-0.123	-0.012	-0.020	-0.000
220425-0603.4	0.7273	-0.062	-0.139	-0.034	-0.027	0.021
181437-1324.0	1.3992	-0.001	-0.110	-0.004	-0.054	0.004
125028-6039.8	2.0439	-0.030	-0.085	-0.010	-0.031	-0.001
162959-4515.5	1.1376	-0.068	-0.155	-0.032	-0.049	0.001
072345-6500.6	3.1240	-0.006	-0.066	-0.003	-0.060	-0.001
185503-3359.5	1.1678	-0.019	-0.070	-0.015	-0.060	0.002
175924-0954.7	2.3438	-0.115	-0.119	-0.090	-0.088	0.006
123140-6237.3	1.9307	-0.009	-0.184	-0.004	-0.051	0.001
172116-3759.2	5.2056	-0.021	-0.135	-0.005	-0.033	0.007
012917-7243.4	180.60	-0.050	-0.190	0.016	-0.052	0.041
051700-5555.4	0.7904	-0.059	-0.126	-0.028	-0.021	0.020
105554-4349.6	1.1541	-0.034	-0.185	-0.019	-0.030	0.037
132831-4107.9	16.515	-0.010	-0.069	-0.011	0.003	-0.015

in compressed files: `ec.tgz` (49MB), `esd.tgz` (27MB), `ed.tgz` (27MB).

Some investigators (e.g. P. Eggleton) strongly prefer classification of close binaries into EW, EB, and EA types, on the grounds that photometry alone cannot provide unique geometry. Our classification into EC, ESD, and ED types should be considered preliminary, as we have only single band photometry and no spectroscopic information for thousands of our binaries.



**Figure 5.** Examples of light curves of 9 contact (EC), 3 semi-detached (ESD) and 3 detached binaries (ED) are shown. All these are ASAS discoveries, even though all of them are very bright. Notice a distinct difference in the light curve shapes between contact and a semi-detached binaries. The first three rows are sorted by minimum depth difference, the fourth row presents contact binaries with long end exceptionally long periods, and the fifth with the distinct maximum height difference. See Table 1 for details.

A sample of the catalog of ASAS eclipsing binaries is shown in Table 2. The full table has the information for all 11,099 binaries, and can be found on our web page, in the file `Vars.E`.

A subset of all ASAS results related to binary stars was ‘frozen’ for the epoch of this paper. ASAS is an on-going project, with more data added every clear night. Therefore, as time goes on, the volume of data related to eclipsing binaries will increase, and the quality will improve. To make the results presented in this paper reproducible we decided to provide ‘frozen’ data.

Note: all light curves presented in Fig. 5 are for bright variables discovered with ASAS survey. They were not known before. Just as expected, the majority of ASAS variables are new discoveries (Paczynski 1997).

#### 4 STATISTICS OF CONTACT BINARIES

All the statistics of this section is based on data provided in the file `Vars.E`, which can be found on our web page. Table

2 is a sample of this file. In particular, the classification of ASAS binaries is given there.


The Galactic distribution of ASAS eclipsing binaries shows a significant difference, with detached binaries most strongly concentrated to the galactic plane, while the short period contact binaries have almost isotropic distribution. It is well known that there is a period - luminosity relation for W UMa stars (Rucinski 1996, Klagyivik and Csizmadia 2004), and the Galactic distribution of those stars is not surprising. The strong concentration of detached binaries to the Galactic plane implies these are even more luminous and massive, on average.

The binary period distribution is shown in Fig. 6 for stars with the Galactic latitude  $|b| > 30^\circ$  to reduce the luminosity bias. The distribution of contact systems (EC) peaks near 0.37 days, it has a sharp cut-off at 0.2 days, and a long tail extending beyond 1 day. In fact the tail extends to more than one hundred days, as it is apparent in Fig. 8 and Fig. 10. Also shown is the distribution of orbital periods of semi-detached (ESD) and detached (ED) binaries.

The important topic of this paper is the presentation

**Table 2.** A sample of the catalog of ASAS eclipsing binaries is presented here. Each star has its own ASAS designation (ID), equatorial coordinates (RA and DEC), orbital period, epoch of minimum light (T0), and ASAS classification. Additional information of other designation and classification is also given, when available. Full Table 2 is available electronically as the file **Vars.E** in our web page.

ID -	RA (2000)	DEC (2000)	Period [days]	T0 2450000+	V [mag]	Amp [mag]	Class -	Other ID	Other Class
065819+1028.4	06:58:19	+10:28:24	0.447763	2387.63	11.04	0.28	EC		
070017+0202.2	07:00:17	+02:02:12	1.397550	1873.07	10.53	0.57	EC	V0460 Mon	EB/KE
211201-2003.2	21:12:01	-20:03:12	0.352202	1872.84	13.22	0.34	EC/ESD		
025535-0219.9	02:55:35	-02:19:54	0.79274	1920.00	11.59	0.54	ESD		
095930-6440.1	09:59:30	-64:40:06	2.2428	1869.42	11.48	1.12	ESD		
202110-4333.9	20:21:10	-43:33:54	1.00218	1876.053	12.08	0.71	ESD/ED	V2265 Sgr	EA
001855-7954.9	00:18:55	-79:54:54	0.90310	1870.45	11.28	0.77	ED		
114500-7247.6	11:45:00	-72:47:36	22.23037	1895.60	11.15	0.09	ED		
235052-2316.7	23:50:52	-23:16:42	1.4023	1871.06	9.42	0.29	ED		



**Figure 6.** The distribution of periods of ASAS contact binaries (EC) at high Galactic latitude  $|b| > 30^\circ$  is plotted. The distribution peaks near 0.37 days, it has a sharp cut-off at 0.2 days, and a tail extending far beyond 1 day. Also shown is the distribution of orbital periods of semi-detached (ESD) and detached (ED) binaries. Contact binaries outnumber other binaries for binary periods shorter than 1 day.


of eclipse depth of EC binaries, as this is directly related to the issue of relaxation oscillations. For contact binaries the depth of both eclipses is directly related to the Fourier coefficients of the light curve. The following analytical formulae approximate these relations with better than 5% accuracy for the majority of stars:

$$D_p = -2(1.2a_2 - 2a_2^2 + (a_1 + a_3)) \quad [\text{mag}], \quad (1)$$

$$D_s = -2(1.2a_2 - 2a_2^2 - (a_1 + a_3)) \quad [\text{mag}], \quad (2)$$

where  $D_p$  and  $D_s$  correspond to the depth of the primary and secondary minimum, respectively. Three examples are presented in Fig. 7.


The distribution of primary eclipse depth with binary period is shown in Fig. 8. A histogram of primary eclipse depth integrated over binary period is shown in Fig. 9. Note that while the vast majority of W UMa binaries have periods




**Figure 7.** The first row of EC stars from Fig. 5 is shown. Vertical lines represent depth of a primary ( $D_p$ ) and a secondary ( $D_s$ ) eclipse calculated with equations (1) and (2), respectively. Flat bottom eclipses may result in a significant error, which is somewhat reduced in the  $D_s/D_p$  ratio.

in the range  $0.2 < P < 1.2$  days, there are contact binaries with periods over 100 days, as it is apparent in Fig. 5 and Fig. 8.


The eclipse depth ratio is shown in Fig. 10 for EC binaries, and in Fig. 11 for EC and ESD binaries. These are important figures. They demonstrate that some EC binaries are in good thermal contact, as the depth of secondary eclipse is almost the same as the primary. But many other EC binaries have the two eclipses which are very unequal, implying that they are not in thermal contact, yet their geometry is close to that expected for a contact system. Finally, there are ESD binaries, in which the geometry is very



**Figure 8.** The distribution of primary eclipse depth as a function of orbital period for contact binaries.  $D_p$  is the fraction of light obscured in the primary eclipse.




**Figure 10.** The ratio of eclipse depth is shown for contact binaries as a function of their period. Stars in thermal contact have the eclipse depth ratio close to one.



**Figure 9.** The distribution of primary eclipse depth integrated over all periods.


different, and of course the two eclipses have usually different depth. Qualitatively, this is just what was expected in the models with relaxation oscillations (cf. Lucy 1976, Flannery 1976, Robertson and Eggleton 1977, Yakut and Eggleton 2005). There is no shortage of stars without thermal contact. Combining ASAS data with models of relaxation oscillations should provide quantitative verification of the theory, but this task is beyond the scope of our paper.

Contact binaries often have maxima of different height,



**Figure 11.** The ratio of the eclipse depths, integrated over binary periods, is shown for contact (EC) and semi-detached (ESD) systems. Stars in a thermal contact have the eclipse depth ratio close to one. Note the step in the histogram at  $D_s/D_p \approx 0.7$ . The same feature is also apparent in Fig. 10 at  $\log P \approx -0.4$ .

with the maximum following the primary (i.e. deeper) eclipse being either higher or lower than the maximum following the secondary eclipse. The distribution of the difference is shown in Fig. 12. The positive value of the  $b1/a2$  parameter indicates that the maximum following the primary eclipse is brighter of the two. The asymmetry in distribution,



**Figure 12.** The distribution of light curve asymmetry of contact binaries is shown - this is known as O’Connell (1951) effect. Most stars have their light maxima of approximately equal height, but there is an asymmetry in the distribution: the maxima that follow the primary minimum are on average higher than the maxima preceding the primary eclipse. The O’Connell effect is likely a consequence of gas streams in these binaries.

i.e. the excess of positive values of  $b1/|a2|$ , is known as the O’Connell (1951) effect (cf. also Ruciński 1997b): the maximum following the primary eclipse is on average brighter, presumably due to stream of gas flowing between the two stars. An extreme case of this phenomenon is shown in V361 Lyr (Kaluzny 1991).

## 5 DISCUSSION

Our conclusion, based on the distribution of eclipse depths (Fig. 10, 11), is that relaxation oscillations, first proposed by Lucy and Flannery, are real. There is no shortage of binaries corresponding to no thermal contact, with a very different depth of the two eclipses. Model calculations of the type recently done by Yakut and Eggleton (2005), when combined with ASAS data, should allow a quantitative verification of the theory.

A very large number of contact or near contact systems reveal features never noticed before: the distribution of eclipse depth ratios as shown in Fig. 10 and Fig. 11 has a distinct break around  $\log P \approx -0.4$  and  $D_s/D_p \approx 0.7$ . We do not speculate on the origin of this feature, but we bring this break to the attention of our readers. This break is best seen in systems with the most robust classification.

The most surprising result of this paper is presented in Fig. 6. A traditional view for the origin of W UMa contact binaries is to assume that they come from detached binaries of comparable periods. For the first time we have approximately complete statistics of binaries of all types with orbital periods shorter than one day, and there are very few detached binaries. Obviously, they are more difficult to find

than either contact or semi - detached systems. As times goes on the statistics of ASAS detached binaries will improve, and a statistical analysis will tell us if there is a problem with the origin of W UMa stars. At this time the contact systems seem to appear ”out of nowhere”.

In about a year we shall have I-band data for our eclipsing binaries. This will make it possible to be more quantitative about the distribution properties, including the distribution in our Galaxy. Also, it will be much easier to quantify the impression about a shortage of detached binaries and the space density of systems of different types: EC, ESD, and ED. At this time it is premature for us to speculate about the outcome of binary statistics while we wait for the I-band data.

While this is an observational paper, written to promote the usefulness of the ASAS catalog of variable stars, we are tempted to speculate about possible interpretation of Fig. 6, which was so surprising to us. We are not consistent with the previous paragraph, but the temptation is hard to resist. The following is a speculative hypothesis.

There has been a gradual emergence of the notion that contact systems have companions (Ruciński and Kaluzny 1982, Chambliss 1992, Hendry and Mochnacki 1998, also Tokovinin 2004). Recently, Pribulla and Ruciński (2005) found that up to 50% of W UMa binaries have companions. This opens up a possibility that Kozai (1962) cycle operates in some such triples, as suggested by Kiseleva, Eggleton, and Mikkola (1998), and by Eggleton and Kiseleva-Eggleton (2001). The mechanical three-body orbital evolution, with a large range of inclinations of the two orbits, inner and outer, may induce large variations in the eccentricity of the inner binary. In time the eccentricity may become large enough to make a contact system out of the inner binary.

In this scenario the inner binary has the initial period that is relatively long, and in most cases no eclipses would be detectable. During a Kozai cycle the inner binary occasionally has a chance to reach a physical contact: it may either merge forming a single star, or it may become a W UMa star. This process is somewhat similar to the model of formation of close binaries in globular clusters (Pooley et al. 2003, and references therein).

The surprisingly small number of short period detached eclipsing binaries as seen in Fig. 6 may indicate that the Kozai cycle is not just a curiosity, but it may be an important channel for forming W UMa stars. This possibility cannot be rejected without careful analysis. After all W UMa stars are rare, with the local space density of just 0.2% of the main sequence stars (Rucinski 2002, and references therein).

## ACKNOWLEDGEMENTS

We are very grateful to P. P. Eggleton, D. Fabrycky, S. Ruciński and K. Stepień for many helpful discussions.

The ASAS project was supported to a large extent by the generous donation of William Golden. This work was partially supported by the Polish MNiI grant no. 1P03D-008-27.

## REFERENCES

Chambliss, C. R. 1992, PASP, 104, 663

- Eggleton, P. P., Kiseleva, L. 2001, ApJ, 562, 1012  
 Flannery, B. P. 1976, ApJ, 205, 217  
 Hazlehurst, J. 1970, MNRAS, 149, 129  
 Hendry, P. D., Mochnacki, S. W. 1998, ApJ, 504, 978  
 Kaluzny, J. 1991, AcA, 41, 17  
 Klagyivik, P., and Csizmadia, Sz. 2004, PADEU, 14, 303  
 Kiseleva, L. G., Eggleton, P. P., Mikkola, 1998, MNRAS, 300, 292  
 Kozai, Y. 1962, AJ, 67, 579  
 Lucy, L. B. 1968a, ApJ, 151, 1123  
 Lucy, L. B. 1968b, ApJ, 153, 877  
 Lucy, L. B. 1976, ApJ, 205, 208  
 O'Connell, D. J. K. 1951, Riverview Pub. 2, 85  
 Paczyński, B. 1997, vsar.conf, 357  
 Pojmański, G. 1997, AcA, 47, 467  
 Pojmański, G. 1998, AcA, 48, 35  
 Pojmański, G. 2000, AcA, 50, 177  
 Pojmański, G. 2002, AcA, 52, 397  
 Pojmański, G. 2003, AcA, 53, 341  
 Pojmański, G., and Maciejewski, G. 2004, AcA, 54, 153  
 Pojmański, G., and Maciejewski, G. 2005, AcA, 55, 97  
 Pojmański, G., Pilecki, B., Szczygiel, D. 2005, astro-ph/050817  
 Pooley, D. et al. 2003, ApJL, 591, L131  
 Pribulla, T., Rucinski, S. M. 2005, private communication  
 Quin, S. 2003, MNRAS, 342, 1260  
 Robertson, J. A., and Eggleton, P. P. 1977, MNRAS, 179, 359  
 Rucinski, S. M. 1996, ASPC, 90, 270  
 Rucinski, S. M. 1997a, AJ, 113, 407  
 Rucinski, S. M. 1997b, AJ, 113, 1112  
 Rucinski, S. M. 1998, AJ, 115, 1135  
 Rucinski, S. 2002, PASP, 114, 1124  
 Rucinski, S. M., Kaluzny, J. 1982, Ap&SS, 88, 433  
 Stepień, K. 2003, IAUS 219, 134  
 Stepień, K. 2005, astro-ph/0510464, submitted to A&A  
 Tokovinin, A. 2004, RMxAC, 21, 7  
 Webbink, R. F. 2003, ASPC, 293, 76  
 Yakut, K., Eggleton, P. P. 2005, ApJ, 629, 1055

This paper has been typeset from a  $\text{\TeX}$ /  $\text{\LaTeX}$  file prepared by the author.